VOL. 632 NOS. 1+2 JOURNAL OF CHROMATOGRAPHY FEBRUARY 19, 1993

CONTENTS

16TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, BALTIMORE, MD, JUNE 14–19, 1992, PART II

Applications: proteins and their constituents (continued)

Isolation of plasma proteins from the clotting cascade by heparin affinity chromatography
by D. Josić, F. Bal and H. Schwinn (Vienna, Austria) 1

High-performance liquid chromatography of amino acids, peptides and proteins. CXXII. Application of experimentally derived retention coefficients to the prediction of peptide retention times: studies with myohemerythrin
by M. C. J. Wilce, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 11

High-performance liquid chromatographic analysis of carbohydrate mass composition in glycoproteins
by M. Kunitani and L. Kresin (Emmeryville, CA, USA) 19

Conformational changes of brain-derived neurotrophic factor during reversed-phase high-performance liquid chromatography
by R. Rosenfeld and K. Benedek (Thousand Oaks, CA, USA) 29

Immunochromatographic analysis of proteins. Identification, characterization and purity determination
by A. Riggin and J. R. Sportsman (Indianapolis, IN, USA) and F. E. Regnier (West Lafayette, IN, USA) 37

Structural characterization of glycoprotein digests by microcolumn liquid chromatography–ionspray tandem mass spectrometry
by J. Liu, K. J. Volk, E. H. Kerns, S. E. Klohr, M. S. Lee and I. E. Rosenberg (Wallingford, CT, USA) 45

High-performance liquid chromatography of tryptophan and other amino acids in hydrochloric acid hydrolysates
by I. Molnár-Perl, M. Pintér-Szakács and M. Khalifa (Budapest, Hungary) 57

Biochemical applications

High-performance liquid chromatographic quantitation of phenylthiocarbamyl muramic acid and glucosamine from bacterial cell walls
by S. R. Hagen (Hamilton, MT, USA) 63

Detection of alkaloids in foods with a multi-detector high-performance liquid chromatographic system
by L. A. Lin (Cincinnati, OH, USA) 69

Comparison of columns of chemically modified porous glass and silica in reversed-phase high-performance liquid chromatography of ginsenosides
by H. Kanazawa, Y. Nagata, E. Kurosaki, Y. Matsushima and N. Takai (Tokyo, Japan) 79

Carbohydrate analysis of fermentation broth by high-performance liquid chromatography utilizing solid-phase extraction
by R. G. Bell and K. L. Newman (Chicago Heights, IL, USA) 87

High-performance liquid chromatography with ion pairing and electrochemical detection for the determination of the stability of two forms of vitamin C
by K. M. De Antonis, P. R. Brown and Z. Yi (Kingston, RI, USA) and P. D. Maugle (Norwich, CT, USA) 91

Estimation of the extent of lipid peroxidation in the ischemic and reperfused heart by monitoring lipid metabolic products with the aid of high-performance liquid chromatography
by G. A. Cordis and N. Maulik (Farmington, CT, USA), D. Bagchi (Omaha, NE, USA) and R. M. Engelman and D. K. Das (Farmington, CT, USA) 97

Size-exclusion chromatographic determination of β-glucan with postcolumn reaction detection
by T. Suortti (Espoo, Finland) 105

Identification and determination of phenolic constituents in natural beverages and plant extracts by means of a coulometric electrode array system
by G. Achilli, G. P. Cellerino and P. H. Gamache (Bedford, MA, USA) and G. V. Melzi d'Eril (Pavia, Italy) 111

Supercritical fluid extraction of synthetic organochlorine compounds in submerged aquatic plants
by P. R. McEachern and G. D. Foster (Fairfax, VA, USA) 119

Electrophoresis

Separation of metallothionein isoforms by capillary zone electrophoresis
by J. H. Beattie (Aberdeen, UK), M. P. Richards (Beltsville, MD, USA) and R. Self (Aberdeen, UK) 127

Temperature-programmed capillary affinity gel electrophoresis for the sensitive base-specific separation of oligodeoxynucleotides
by Y. Baba and M. Tsuhako (Kobe, Japan) and T. Sawa and M. Akashi (Kagoshima, Japan) 137

Determination of ten β-blockers in urine by micellar electrokinetic capillary chromatography
by P. Lukkari, H. Sirén, M. Pantsar and M.-L. Riekkola (Helsinki, Finland) 143

Voltage programming in capillary zone electrophoresis
by H.-T. Chang and E. S. Yeung (Ames, IA, USA) 149

Fundamental and practical aspects of coupled capillaries for the control of electroosmotic flow in capillary zone electrophoresis of proteins
by W. Nashabeh and Z. El Rassi (Stillwater, OK, USA) 157

Determination of sulphonamides in pharmaceuticals by capillary electrophoresis
by C. L. Ng, H. K. Lee and S. F. Y. Li (Singapore, Singapore) 165

Capillary sodium dodecyl sulfate gel electrophoresis of proteins
by A. Guttman, J. A. Noland and N. Cooke (Fullerton, CA, USA) 171

Prediction of the migration behavior of organic acids in micellar electrokinetic chromatography
by S. C. Smith and M. G. Khaledi (Raleigh, NC, USA) 177

Polymer- and surfactant-coated capillaries for isoelectric focusing
by X.-W. Yao and F. E. Regnier (West Lafayette, IN, USA) 185

Capillary electrophoresis of glycoconjugates in alkaline media
by M. Stefansson and D. Westerlund (Uppsala, Sweden) 195

Observation of flow profiles in electroosmosis in a rectangular capillary
by T. Tsuda and M. Ikedo (Nagoya, Japan) and G. Jones, R. Dadoo and R. N. Zare (Stanford, CA, USA) 201

On-line sample preconcentration on a packed-inlet capillary for improving the sensitivity of capillary electrophoretic analysis of pharmaceuticals
by M. E. Swartz and M. Merion (Milford, MA, USA) 209

OTHER SEPARATION METHODS

Screening of β-blockers in human serum by ion-pair chromatography and their identification as methyl or acetyl derivatives by gas chromatography–mass spectrometry
by H. Sirén, M. Saarinen, S. Hainari, P. Lukkari and M.-L. Riekkola (Helsinki, Finland) 215

Isoelectric focusing field-flow fractionation and capillary isoelectric focusing with electroosmotic zone displacement. Two approaches to protein analysis in flowing streams
by J. Chmelík (Brno, Czechoslovakia) and W. Thormann (Berne, Switzerland) 229

Author Index Vols. 631 and 632 235